Mon. Not. R. Astron. Soc. **000**, 000–000 (0000) Printed 24 April 2013 (MN LaTeX style file v2.2)

arXiv:1304.6120v1 [astro-ph.CO] 22 Apr 2013

Green's function of the cosmological thermalization problem

J. Chluba[1]⋆
[1] *Johns Hopkins University, Department of Physics and Astronomy, Bloomberg Center 435, 3400 N. Charles St., Baltimore, MD 21218*

Accepted 2013 April. Received 2013 April 10

ABSTRACT
Energy release in the early Universe leads to spectral distortions of the cosmic microwave background (CMB) which in the future might allow probing different physical processes in the pre-recombination ($z \gtrsim 10^3$) epoch. Depending on the energy injection history, the associated distortion partially thermalizes due to the combined action of Compton scattering, double Compton scattering and Bremsstrahlung emission, a problem that in general is hard to solve. Various analytic approximations describing the resulting distortion exist, however, for small distortions and fixed background cosmology the Green's function of the problem can be computed numerically. Here we show that this approach gives very accurate results for general thermal histories, allowing fast and quasi-exact computation of the spectral distortion given the energy release rate. Our method can thus be used to forecast possible constraints on early-universe physics obtained from future measurements of the CMB spectrum.

Key words: Cosmology: cosmic microwave background – theory – observations

1 INTRODUCTION

The observations of the cosmic microwave background (CMB) with COBE/FIRAS (Mather et al. 1994; Fixsen et al. 1996) showed that the CMB frequency spectrum is extremely close to a perfect blackbody, allowing deviations of no more than $\Delta I_\nu / I_\nu \lesssim$ few $\times 10^{-5}$ at wavelengths ranging from $100\,\mu$m to 1 cm. Dramatic improvements in experimental design and technology over the past decades since the launch of COBE may soon allow much more sensitive measurements of the CMB frequency spectrum (e.g., see Fixsen & Mather 2002; Kogut et al. 2011). For this reason, CMB spectral distortion have recently experienced renewed interest (e.g., see Chluba & Sunyaev 2012; Chluba et al. 2012b; Pajer & Zaldarriaga 2012; Khatri & Sunyaev 2012b; Tashiro et al. 2012; Ali-Haïmoud 2013; Sunyaev & Khatri 2013, to name a few), potentially providing a way to probe new physics. It therefore is time to ask what exactly one might hope to learn about the thermal history of our Universe from future measurements of the CMB spectrum and how the information can be extracted.

To address this problem, a fast and accurate method for computing the spectral distortion from different scenarios is required. Previous numerical approaches (e.g., see Procopio & Burigana 2009; Chluba & Sunyaev 2012) are rather time-consuming. Simple analytic approximations exist (e.g., see Sunyaev & Zeldovich 1970b; Illarionov & Sunyaev 1974; Burigana et al. 1991; Hu & Silk 1993; Burigana et al. 1995; Khatri & Sunyaev 2012b,a), however, for small distortions and fixed background cosmology it is possible to obtain a numerical approximation for the Green's function of the thermalization problem. This allows us to compute the spectral distortion for general thermal histories in an economic and quasi-exact way. In this short paper, we present our method which provides the basis for detailed forecasts, assessing the future experimental possibilities for spectral distortion measurements, and comparisons with independent (Khatri & Sunyaev 2013) approaches.

2 COSMOLOGICAL THERMALIZATION PROBLEM

In the early Universe, photons undergo many interactions with the electrons and baryons. A detailed discussion of the cosmological thermalization problem for different energy-release mechanisms can be found in Chluba & Sunyaev (2012). Compton scattering allows photons to exchange energy with the electrons and baryons, causing diffusion in frequency, while double Compton and Bremsstrahlung emission are responsible for adjusting the photon number. The Boltzmann equation for the evolution of the average photon occupation number, $n_\nu(t)$, can be expressed as:

$$\frac{\partial n_\nu}{\partial t} - H\,\nu\,\frac{\partial n_\nu}{\partial \nu} = \left.\frac{\mathrm{d}n_\nu}{\mathrm{d}t}\right|_{\rm C} + \left.\frac{\mathrm{d}n_\nu}{\mathrm{d}t}\right|_{\rm DC} + \left.\frac{\mathrm{d}n_\nu}{\mathrm{d}t}\right|_{\rm BR} + \left.\frac{\mathrm{d}n_\nu}{\mathrm{d}t}\right|_{\rm S}. \quad (1)$$

The second term on the left hand side describes the redshifting of photons due to the adiabatic expansion of the Universe and the right hand side terms correspond to the physical processes quoted above. We also explicitly added a photon source term, $\mathrm{d}n_\nu/\,\mathrm{d}t|_{\rm S}$, however, direct photon production is normally negligible. If, for example, energy release is caused by some decaying or annihilating relic particle, very few soft photons are directly produced, and high-energy photons and decay products just heat the medium (e.g., see Shull & van Steenberg 1985; Chen & Kamionkowski 2004; Hütsi et al. 2009; Slatyer et al. 2009; Cirelli et al. 2009). Subsequently, this leads to up-scattering of CMB photons and thus

---

⋆ E-mail:jchluba@pha.jhu.edu

© 0000 RAS

sources distortions, initiating the cosmological thermalization process

In addition to the photon occupation number, one has to follow the temperature of ordinary matter, including the cooling/heating due to the adiabatic expansion of the Universe, Compton scattering, double Compton emission and Bremsstrahlung. Extra energy release can be parametrized using an effective heating rate, $\dot{Q}(t)$. It increases the matter temperature, subsequently causing up-scattering of CMB photons by Compton scattering and photon production at very low frequencies (Sunyaev & Zeldovich 1970a).

For us it is important that any primordial spectral distortion has to be *tiny* (cf., Fixsen et al. 1996; Fixsen 2009), so that the thermalization problem can be linearized[1]. We furthermore know the main cosmological parameters, such as the baryon density or Hubble expansion rate, extremely well from measurements of the CMB anisotropies (Bennett et al. 2003; Planck Collaboration et al. 2013). Thus it is possible to numerically compute the Green's function, $G_{\rm th}(\nu, z_{\rm h}, z_{\rm obs})$, of the thermalization problem using CosmoTherm. Here $z_{\rm h}$ denotes the heating redshift and $z_{\rm obs}$ the observing redshift. The resulting spectral distortion at $z = 0$ is then given by

$$\Delta I_\nu(z=0) = \int G_{\rm th}(\nu, z', 0)\,\frac{{\rm d}(Q/\rho_\gamma)}{{\rm d}z'}\,{\rm d}z', \qquad (2)$$

where $\rho_\gamma(z)$ is the energy density of the CMB photon field, and we introduced the intensity, $\Delta I_\nu \equiv (2h\nu^3/c^2)\Delta n_\nu$. We chose to use redshift as time variable, since most energy release mechanisms have rather simple form in this coordinate (see Chluba & Sunyaev 2012). From the practical point of view this poses no significant limitation, and Eq. (2) can be generalized to other variables, as long as the heating rate varies slowly.

3 RESULTS FOR THE GREEN'S FUNCTION

To obtain the Green's function we solve the thermalization problem assuming that the heating rate is given by a δ-function centered at heating redshift, $z_{\rm h}$. Numerically we simply approximate the heating rate by a narrow Gaussian in redshift

$$\left.\frac{{\rm d}(Q/\rho_\gamma)}{{\rm d}z}\right|_\delta \approx \frac{\Delta\rho_{\gamma,{\rm h}}}{\rho_{\gamma,{\rm h}}}\,\frac{\exp\left(-[z-z_{\rm h}]^2/2\sigma_z^2\right)}{\sqrt{2\pi\sigma_z^2}} \qquad (3)$$

with[2] $\sigma_z \simeq 0.01$ and $\Delta\rho_{\gamma,{\rm h}}/\rho_{\gamma,{\rm h}} \simeq 10^{-6}$. By evolving the distortion, Δn_ν, and matter temperature, $T_{\rm m}$, from $z_{\rm s} \gg z_{\rm h}$ until late times ($z \simeq 200$) we obtain an approximation for the Green's function from $G_{\rm th}(\nu, z_{\rm h}, 0) \approx \Delta n_\nu^{\rm num}/[\Delta\rho_{\gamma,{\rm h}}/\rho_{\gamma,{\rm h}}]$. At $z_{\rm s}$ the photon field is given by a blackbody at temperature $T_{\rm s} < T_0(1+z_{\rm s})$, where $T_0 = 2.726\,{\rm K}$ (Fixsen 2009) is the CMB temperature today. Initially, also the temperature of ordinary matter is $T_{\rm m}(z_{\rm s}) \equiv T_{\rm s}$. At the end of the evolution, we ensure that the effective photon temperature is close to T_0. To compute the Green's function we furthermore exclude the down-scattering of photons by adiabatically cooling electrons, leading to a distortion characterized by *negative* chemical potential and y-parameter (Chluba 2005; Chluba & Sunyaev 2012; Khatri et al. 2012). This process can be included afterward, since for fixed cosmology the corresponding distortion is known precisely. We adopted the following cosmological parameters: $Y_{\rm p} = 0.24$, $\Omega_{\rm m} = 0.26$, $\Omega_{\rm b} = 0.044$, $\Omega_\Lambda = 0.74$, $\Omega_{\rm k} = 0$, $h = 0.71$, and $N_{\rm eff} = 3.046$, however, since the computation is rather straightforward, it is very easy to adjust these values.

3.1 Numerical results

In Fig. 1 we present the numerical results for the Green's function of the cosmological thermalization problem. For very early energy release ($z_{\rm h} \gg z_\mu \simeq 2\times10^6$), it is equivalent to a temperature shift, having the shape

$$4G_{\rm th}(\nu, z_{\rm h} \gg z_\mu, 0) \approx G(\nu) = \frac{2h\nu^3}{c^2}\,\nu\partial_\nu n^{\rm pl}(\nu, T_0) = \frac{2h\nu^3}{c^2}\,\frac{x\,{\rm e}^x}{({\rm e}^x-1)^2},$$

where $x = h\nu/kT_0$, $n^{\rm pl}(\nu, T_0) = 1/({\rm e}^x - 1)$, and the factor of 4 arises from $\Delta\rho_\gamma/\rho_\gamma \simeq 4\Delta T/T$. This indicates that the thermalization process is extremely efficient, with the residual μ-type spectral distortion (Sunyaev & Zeldovich 1970a) being exponentially suppressed by the distortion visibility function[3], $\mathcal{J}(z_{\rm h}) = {\rm e}^{-(z_{\rm h}/z_\mu)^{5/2}}$ (Danese & de Zotti 1982; Hu & Silk 1993). To improve the accuracy of our computation, at $z \gtrsim 5\times10^4$ we treated the temperature shift independently, imposing photon number conservation, $\int \nu^2 \Delta n_\nu\,{\rm d}\nu \equiv 0$, to define the distortion part[4]. Since the temperature shift caused by early energy release is not directly observable, this procedure allowed us to obtain precise results for the μ-type distortion related to significant energy release at very early times.

For low heating redshift ($z_{\rm h} \lesssim 10^4$), the Green's function takes the shape of a Compton y-distortion (Zeldovich & Sunyaev 1969). At this redshift, the effective y-parameter remains very small, so that photons only weakly interact with the electrons and the distortion is described by

$$4G_{\rm th}(\nu, z_{\rm h} \lesssim 10^4, 0) \approx Y_{\rm SZ}(\nu) = \frac{2h\nu^3}{c^2}\,\frac{x\,{\rm e}^x}{({\rm e}^x-1)^2}\,[x\coth(x/2) - 4]\,.$$

At very low frequencies ($\nu \lesssim 1\,{\rm GHz}$) this approximation is not as accurate, since matter and radiation start decoupling and Bremsstrahlung emission (or absorption) can alter the spectrum (see Chluba & Sunyaev 2012, for illustration). However, the total number of photons in that part of the radiation field is very small and this effect will be challenging to observe. The numerical results for the Green's function include these aspects.

At $z_{\rm h} \simeq 3\times10^5$ the Green's function is mainly represented by a μ-type distortion, obtained using the condition $\int \nu^2\Delta n_\nu\,{\rm d}\nu \equiv 0$ (see Chluba et al. 2012b; Khatri & Sunyaev 2012a, for more detail):

$$\alpha^{-1}G_{\rm th}(\nu, z_{\rm h} \simeq 3\times10^5, 0) \approx M(\nu) = \frac{2h\nu^3}{c^2}\,\frac{{\rm e}^x}{({\rm e}^x-1)^2}\,[x/\beta - 1]\,.$$

Here $\alpha \approx 1.401$ (Sunyaev & Zeldovich 1970a; Illarionov & Sunyaev 1974) and $\beta = 3\zeta(3)/\zeta(2) \approx 2.1923$. Neglecting the temperature shift, the Green's function for the distortion part is well represented by $M(\nu)$, even for $z_{\rm h} \gtrsim 3\times10^5$ when additionally multiplying with the distortion visibility function, $\mathcal{J}(z_{\rm h})$, giving the approximation $G_{\rm th}(\nu, z_{\rm h} \gtrsim 3\times10^5, 0) \approx 1.401\mathcal{J}(z_{\rm h})\,M(\nu) + \frac{1-\mathcal{J}(z_{\rm h})}{4}\,G(\nu)$.

---

[1] We numerically solve the full non-linear system, however, limiting the total energy release to $\Delta\rho_\gamma/\rho_\gamma \ll 1$.

[2] We confirmed the convergence of the Green's function by changing the width of the Gaussian, finding excellent agreement between the results.

[3] See Khatri & Sunyaev (2012b) for recent analytical discussion.

[4] The thermalization process slowly increases the photon number density or equivalently the effective temperature (defined by $N_\gamma \propto T^3$). To avoid having a *large* contribution from $G(\nu)$ to the numerical solution we can restart the computation at different stages, shifting the temperature of the reference blackbody to the new effective temperature of the photon field. At $z \lesssim 5\times10^4$ thermalization is already very inefficient and no significant *average* temperature shift can be produced, so that we leave the temperature of the reference blackbody unchanged.



Figure 1. Numerical results for the Green's function of the cosmological thermalization problem for various heating redshifts, $z_{\rm h} \in [10^3, 5\times10^6]$. Energy release at very high redshifts causes an increase in the effective temperature of the photon field, while at low redshifts photons only partially up-scatter, creating a y-distortion. Around $z_{\rm h} \simeq 3\times10^5$ a pure μ-distortion is created. All intermediate stages are roughly (precision $\simeq$ 10% – 30%) represented by a superposition of these extreme cases, however, the small residuals provide a principle possibility to distinguish different thermal histories at redshifts $10^4 \lesssim z \lesssim 3\times10^5$.

Finally, Figure 1 also shows the Green's function at several intermediate stages, $10^4 \lesssim z_{\rm h} \lesssim 3\times10^5$. Signals produced mainly during this epoch were already discussed in Chluba & Sunyaev (2012, see Fig. 15 and 19), indicating that the total distortion is not simply given by a pure superposition of μ and y-distortion. More recently, this was also demonstrated by Khatri & Sunyaev (2012a). The small ($\simeq$ 10% – 30%) residuals might allow distinguishing different energy release scenarios in the future. This is especially interesting if, for instance, decaying particles with lifetimes $t_{\rm X} \simeq 2.6\times10^8\,{\rm sec} - 2.2\times10^{11}\,{\rm sec}$ are present (Chluba & Sunyaev 2012), however, detailed forecasts are required to address this question. One can still obtain a fairly good approximation for the Green's function in this intermediate regime by assuming that the spectrum is close to a superposition of a pure μ- and y-distortion, $G_{\rm th}(\nu, z_{\rm h} \lesssim 3\times10^5, 0) \approx 1.401\mathcal{J}_\mu(z_{\rm h})\,M(\nu) + \frac{\mathcal{J}_y(z_{\rm h})}{4}\,Y_{\rm SZ}(\nu)$. The coefficients, $\mathcal{J}_y(z_{\rm h})$ and $\mathcal{J}_\mu(z_{\rm h})$, which approximate the transition between the μ- and y-distortion regime, can be determined using a least square fit to the Green's function. We find

$$\mathcal{J}_y(z_{\rm h}) \approx \left(1 + \left[\frac{1+z}{6.0\times10^4}\right]^{2.58}\right)^{-1}$$
$$\mathcal{J}_\mu(z_{\rm h}) \approx 1 - \exp\left(-\left[\frac{1+z}{5.8\times10^4}\right]^{1.88}\right). \tag{4}$$

This approximation for $G_{\rm th}(\nu, z_{\rm h}, 0)$ works at $\simeq$ 10% – 30% precision for the standard cosmology, with the residuals containing additional information about the time-dependence of the energy release at $10^4 \lesssim z_{\rm h} \lesssim 3\times10^5$. In summary, this means that a pretty good approximation for the Green's function at all redshifts is given by

$$G^*_{\rm th}(\nu, z_{\rm h}, 0) = 1.401\mathcal{J}_\mu(z_{\rm h})\,\mathcal{J}(z_{\rm h})\,M(\nu) + \frac{\mathcal{J}_y(z_{\rm h})}{4}\,Y_{\rm SZ}(\nu) + \frac{1-\mathcal{J}(z_{\rm h})}{4}\,G(\nu). \tag{5}$$

The redshift-dependence of the functions, $\mathcal{J}_y(z_{\rm h})$, $\mathcal{J}_\mu(z_{\rm h})$ and $\mathcal{J}(z_{\rm h})$ is illustrated in Fig. 2. At $z \simeq 5.3\times10^4$ we find $\mathcal{J}_y(z_{\rm h}) \simeq \mathcal{J}_\mu(z_{\rm h})$, justifying why $z \simeq 5\times10^4$ was previously used to mark the transition between μ- and y-era (Hu & Silk 1993).

It is also important to mention that the energy integral $\int [G^*_{\rm th}(\nu, z_{\rm h}, 0)/\rho_\gamma]\,{\rm d}\nu = \left[\mathcal{J}_\mu(z_{\rm h})\mathcal{J}(z_{\rm h}) + \mathcal{J}_y(z_{\rm h}) + 1 - \mathcal{J}(z_{\rm h})\right] \lesssim 1$. The 'missing' energy can be found in the residual distortion, which is not represented well by the approximation, although it never exceeds $\simeq$ 16% (maximum at $z \simeq 7.7\times10^4$). For simple estimates, Eq. (5) should thus be very useful, while for high precision the exact form of the Green's function must be used.

3.2 Spectral distortion for some examples

We implemented the Green's function method as part of CosmoTherm[5]. With Eq. (2) this allows very fast computation (a few seconds versus $\simeq$ 10 min – 1 h) of the resulting spectral distortion for given thermal history. In Fig. 2 we show the heating rate and resulting distortion for some illustrative cases. We confirmed the precision of the Green's function approach by explicitly computing the distortion for different scenarios solving the evolution equations with CosmoTherm, finding excellent agreement [$\simeq$ 1%-level uncertainties in the background cosmology are more rele-

---

[5] CosmoTherm is available at `www.Chluba.de/CosmoTherm`.



Figure 2. Effective heating rate and spectral distortion created by different scenarios. In all cases the total energy release was similar to $\Delta\rho_\gamma/\rho_\gamma \simeq$ few $\times 10^{-8}$ (irrespective of observational constraints). We only included the distortion part for the calculation since the shift in the overall temperature of the photon field remains unobservable. For comparison we show a y-distortion with $y \simeq 10^{-8}$. We also illustrate the redshift-dependence of $\mathcal{J}_y(z_{\rm h})$, $\mathcal{J}_\mu(z_{\rm h})$ and $\mathcal{J}(z_{\rm h})$, which we multiplied by 10^{-9}, for convenience. The different scenarios are described in more detail in Sect. 3.2.

vant]. The two annihilation scenarios assume different cross sections, representing s-wave [$\langle\sigma v\rangle \simeq$ const], and p-wave annihilation [$\langle\sigma v\rangle \simeq T/m \propto (1+z)$] (see McDonald et al. 2001, for related discussion). Strong constraints for s-wave annihilation of dark matter can be derived from measurements of the CMB temperature anisotropies (Galli et al. 2009; Hütsi et al. 2009, 2011; Planck Collaboration et al. 2013), however, CMB anisotropy limits on the p-wave scenario are much weaker because most of the energy is liberated prior to the recombination epoch ($z \simeq 10^3$). We can see that for p-wave annihilation most of the energy is released during the μ-era, while for s-wave annihilation heating at low redshifts also contributes significantly. Spectral distortions thus might allow shedding light on these cases, although degeneracies with other scenarios are expected.

For illustration we also chose a decaying particle scenario with the main heating occurring at $z \simeq 5 \times 10^4$. As explained above, in this case the distortion is not just given by a simple superposition of (pure) μ and y-distortion, so that this scenarios might be distinguishable from, for example, the signal created by the dissipation of acoustic modes (Chluba et al. 2012b,a), at least when the simplest shape for the primordial power spectrum is assumed. A more detailed forecast will be required[6], and we plan to apply our method, assuming experimental settings similar to PIXIE (Kogut et al. 2011).

3.3 Cosmology-dependence of the Green's function

The cosmology-dependence of the Green's function is mainly driven by (i) the CMB monopole temperature; (ii) the number density of baryons and electrons; (iii) the faction of helium atoms (though the Bremsstrahlung Gaunt-factors and the effect on the number of electrons per baryon); and (iv) the expansion rate of the Universe (i.e., $N_{\rm eff}$ and at later stages $\Omega_{\rm m}$). Since the cosmological parameters are known with percent-level precision, the Green's function is accurate to a similar level. For detailed parameter forecasts the cosmology-dependence of the Green's function could thus lead to significant degeneracies, if differences at the level of a few percent matter. This can be easily taken into account, however, we leave an in depth discussion to future work.

4 CONCLUSIONS

We computed the Green's function for the cosmological thermalization problem assuming standard cosmological parameters. With our method it is possible to predict the resulting spectral distortion signal for different energy release scenarios with high precision at practically no computational cost (see Fig 2 for some examples). This approach is now part of CosmoTherm and will be very useful for detailed forecasts of possible future constraints on the pre-recombination thermal history of our Universe derived from measurements of the CMB spectrum.

In particular, it will be very important to ask how degenerate different scenarios are. For example, energy release related to dark matter annihilation with s-wave cross section is expected to give rise to a distortion that is practically degenerate with those caused by dissipation of adiabatic modes in the early Universe, if the spectral index $n_{\rm S}$ of the primordial power spectrum at wavenumber $k \gtrsim 1\,{\rm Mpc}^{-1}$ is scale-invariant (e.g. see Chluba & Sunyaev 2012). One the other hand, recent constraints from Planck (Planck Collaboration et al. 2013) show that at large scales $n_{\rm S} \simeq 0.96$. Extrapolating this to small scales thus suggests that the shape of the distortion ought be different in the two cases. Our computations also show that thermal histories with decaying relics might be constrainable if the lifetime of the particle is $t_X \simeq 2.6 \times 10^8\,{\rm sec} - 2.2 \times 10^{11}\,{\rm sec}$. We look forward to exploring these possibilities in the future.

ACKNOWLEDGMENTS

JC thanks Yacine Ali-Haïmoud, Rishi Khatri and Rashid Sunyaev for useful discussions and comments on the paper. He also thanks Rajat Thomas for great hospitality during his visit to Amsterdam, where most of this work was finished. Use of the GPC supercomputer at the SciNet HPC Consortium is acknowledged. SciNet is funded by: the Canada Foundation for Innovation under the auspices of Compute Canada; the Government of Ontario; Ontario Research Fund - Research Excellence; and the University of Toronto. JC was support by the grants DoE SC-0008108 and NASA NNX12AE86G.

REFERENCES

Ali-Haïmoud Y., 2013, Phys.Rev.D, 87, 023526
Bennett C. L. et al., 2003, ApJS, 148, 1
Burigana C., Danese L., de Zotti G., 1991, A&A, 246, 49
Burigana C., de Zotti G., Danese L., 1995, A&A, 303, 323
Chen X., Kamionkowski M., 2004, Phys.Rev.D, 70, 043502
Chluba J., 2005, PhD thesis, LMU München
Chluba J., Erickcek A. L., Ben-Dayan I., 2012a, ApJ, 758, 76
Chluba J., Khatri R., Sunyaev R. A., 2012b, MNRAS, 425, 1129
Chluba J., Sunyaev R. A., 2012, MNRAS, 419, 1294
Cirelli M., Iocco F., Panci P., 2009, JCAP, 10, 9
Danese L., de Zotti G., 1982, A&A, 107, 39
Fixsen D. J., 2009, ApJ, 707, 916
Fixsen D. J., Cheng E. S., Gales J. M., Mather J. C., Shafer R. A., Wright E. L., 1996, ApJ, 473, 576
Fixsen D. J., Mather J. C., 2002, ApJ, 581, 817
Galli S., Iocco F., Bertone G., Melchiorri A., 2009, Phys.Rev.D, 80, 023505
Hu W., Silk J., 1993, Phys.Rev.D, 48, 485
Hütsi G., Chluba J., Hektor A., Raidal M., 2011, A&A, 535, A26
Hütsi G., Hektor A., Raidal M., 2009, A&A, 505, 999
Illarionov A. F., Sunyaev R. A., 1974, Astronomicheskii Zhurnal, 51, 1162
Khatri R., Sunyaev R. A., 2012a, JCAP, 9, 16
Khatri R., Sunyaev R. A., 2012b, JCAP, 6, 38
Khatri R., Sunyaev R. A., 2013, ArXiv:1303.7212
Khatri R., Sunyaev R. A., Chluba J., 2012, A&A, 540, A124
Kogut A. et al., 2011, JCAP, 7, 25
Mather J. C. et al., 1994, ApJ, 420, 439
McDonald P., Scherrer R. J., Walker T. P., 2001, Phys.Rev.D, 63, 023001
Pajer E., Zaldarriaga M., 2012, Physical Review Letters, 109, 021302
Planck Collaboration et al., 2013, ArXiv:1303.5076
Procopio P., Burigana C., 2009, A&A, 507, 1243
Shull J. M., van Steenberg M. E., 1985, ApJ, 298, 268
Slatyer T. R., Padmanabhan N., Finkbeiner D. P., 2009, Physical Review D (Particles, Fields, Gravitation, and Cosmology), 80, 043526
Sunyaev R. A., Khatri R., 2013, ArXiv:1302.6553
Sunyaev R. A., Zeldovich Y. B., 1970a, Astrophysics and Space Science, 9, 368
Sunyaev R. A., Zeldovich Y. B., 1970b, Astrophysics and Space Science, 7, 20
Tashiro H., Sabancilar E., Vachaspati T., 2012, Phys.Rev.D, 85, 103522
Zeldovich Y. B., Sunyaev R. A., 1969, Astrophysics and Space Science, 4, 301

---

[6] See Khatri & Sunyaev (2013) for some recent discussion.